## Japan Geoscience Union Meeting 2014

(28 April - 02 May 2014 at Pacifico YOKOHAMA, Kanagawa, Japan)

©2014. Japan Geoscience Union. All Rights Reserved.



PCG38-P04

Room:Poster

Time:May 1 18:15-19:30

## Two dimensional numerical study on Venusian gravity waves by using mesoscale model

ANDO, Hiroki<sup>1\*</sup>; SUGIYAMA, Ko-ichiro<sup>1</sup>; IMAMURA, Takeshi<sup>1</sup>; ODAKA, Masatsugu<sup>2</sup>; NAKAJIMA, Kensuke<sup>3</sup>

<sup>1</sup>ISAS/JAXA, <sup>2</sup>Hokkaido University, <sup>3</sup>Kyusyu University

Recently Venusian gravity waves are often observed. For example, Airglow measurements of  $O_2$  found the gravity waves with horizontal wavelength of  $\sim 100$  km at 110 km altitude. UV images also detect gravity waves with the horizontal wavelength of 60-150 km at the cloud top level (70 km altitude). However, only a specific altitudes can be observed in these measurements, thus it is difficult to examine the propagation characteristics and momentum flux of waves. Radio occultation measurements also detect upward propagating waves from the vertical temperature profiles within the altitude range of 65-90 km and suggests that waves with the vertical wavelength of 5-10 km are dominant by the spectral analysis. However, horizontal resolution in this measurement is  $\sim 200$  km, then small scale gravity waves cannot be observed. Therefore, it is difficult to understand how these gravity waves have their influence on the Venusian atmosphere.

In this study we developed a new Venusian mesoscale model and examined the propagation characteristic of the waves. In the model, we simulated the generation and propagation of the waves including the convective motion in the Venusian cloud layer. We will make a presentation about the initial analysis results.

Keywords: Venus atmosphere, Gravity waves, Numerical study